

Aspects of short range correlations in a relativistic model

P.K. Panda and D.P. Menezes

Depto de Física-CFM, Universidade Federal de Santa Catarina, CP. 476, 88040-900 Florianópolis-SC, Brazil

C. Providência and J. Da Providência

Centro de Física Teórica - Dep. de Física, Universidade de Coimbra, P-3004 - 516 Coimbra, Portugal

In the present work short range correlations are introduced for the first time in a relativistic approach to the equation of state of the infinite nuclear matter in the framework of the Hartree-Fock approximation using an effective Hamiltonian derived from the $\sigma - \omega$ Walecka model. The unitary correlation method is used to introduce short range correlations. The effect of the correlations in the ground state properties of the nuclear matter is discussed.

Nuclear physics is an effective theory of the nucleus regarded as a system of nucleons. In this theory the essential degrees of freedom are the center of mass coordinates of the nucleons as well as their spins and isospins, the interactions being expressed as nucleon-nucleon forces. It is usually accepted, however, that the nucleon-nucleon interaction becomes strongly repulsive at short distances in the relative coordinate of two nuclear particles. It is seen that the phase shifts for $^1S_0$ and $^3S_1$ are positive at low and becomes negative at higher energies [1]. This indicates a repulsive core at short distances and attraction at long distances. There are attempts to derive the nucleon-nucleon force using chiral perturbation theory [2]. In this approach, one- and two-pion exchange contributions are taken into account up to the third chiral order. However, in order to reproduce the nucleon scattering data and the D-wave phase shifts by this method, an *ad hoc* contact interaction (which represents the short range force) must also be included. The realistic nucleon-nucleon forces are basically phenomenological. The Bonn interactions [3] are based on meson exchange which is treated in a relativistic non-local manner. The Argonne interactions [4], on the other hand, describe the pion exchange in a local approximation, the short and medium range nuclear interactions being controlled by phenomenological parameters.

Nonrelativistic calculations based on realistic NN potentials predict equilibrium points that are not able to describe simultaneously the correct binding energy and saturation density; either the saturation density is correct but the binding energy is too small, or the correct binding energy is obtained at a too high density. This behaviour is normally referred to “Coester line” [5]. This problem is generally circumvented through the introduction of a three-body repulsive force [6], or density-dependent repulsive mechanisms. A mechanism of this kind is already present in relativistic models. Due to Lorentz covariance and self-consistency, as the nuclear density increases, the nucleon effective mass decreases. As a result there is a reduction of the attractive force and a net increase of the repulsive force. In fact, the relativistic mean field (RMF) theory formulated by Teller and the others [7] and by Walecka [8] is successful in describing both infinite nuclear matter and finite nuclei. Moreover, the relativistic Hartree-Fock (HF) approximation [9, 10] has also been used for the description of finite nuclei and infinite nuclear matter.

In this paper, we have focused on the treatment of short range correlations in dense nuclear matter for the first time using relativistic equations of motion. Although there are several procedures which may be used to introduce short range correlations into the model wave function, we have preferred to work with the unitary operator method as proposed by Villars [11]. There are several advantages in using an unitary model operator. In particular, one automatically guarantees that the correlated state is normalized. Further, we shall see that for calculation of correlation corrections to the matrix elements of one- and two-body operators is a simple task. The general idea exists for a long time [12, 13, 14] but has not been pursued for the relativistic case. This has been so because in non-relativistic models the interaction arises from the interplay between a long range attraction and a very strong short range repulsion so that, indeed, it is indispensable to take short range correlations into account. In relativistic mean field models, the parameters are phenomenological, fitted to the saturation properties of nuclear matter. Short range correlation effects are clearly included in the model parameters. We expect that the values of these parameters are more ‘fundamental’, or less artificial, when correlations are not neglected.

We start with the effective Hamiltonian as

$$H = \int \psi_\alpha^\dagger(\vec{x})(-i\vec{\alpha} \cdot \vec{\nabla} + \beta M)_{\alpha\beta} \psi_\beta(\vec{x}) d\vec{x} + \frac{1}{2} \int \psi_\alpha^\dagger(\vec{x}) \psi_\gamma^\dagger(\vec{y}) V_{\alpha\beta,\gamma\delta}(|\vec{x} - \vec{y}|) \psi_\delta(\vec{y}) \psi_\beta(\vec{x}) d\vec{x} d\vec{y} \quad (1)$$

with

$$V_{\alpha\beta,\gamma\delta}(r) = (\beta)_{\alpha\beta}(\beta)_{\gamma\delta} V_\sigma(r) + (\delta_{\alpha\beta}\delta_{\gamma\delta} - \vec{\alpha}_{\alpha\beta} \cdot \vec{\alpha}_{\gamma\delta}) V_\omega(r) \quad (2)$$

where

$$V_\sigma(r) = -\frac{g_\sigma^2}{4\pi} \frac{e^{-m_\sigma r}}{r}, \quad V_\omega(r) = \frac{g_\omega^2}{4\pi} \frac{e^{-m_\omega r}}{r}, \quad (3)$$

and $\vec{\alpha}$ are the Dirac-matrices. In the above, ψ is the nucleon field interacting through the scalar and vector potentials. The equal time quantization condition for the nucleons is given by

$$[\psi_\alpha(\vec{x}, t), \psi_\beta(\vec{y}, t)^\dagger]_+ = \delta_{\alpha\beta} \delta(\vec{x} - \vec{y}), \quad (4)$$

where α and β refer to the spin indices. We now also have the field expansion for the nucleons ψ at time $t=0$ given as [15]

$$\psi(\vec{x}) = \frac{1}{\sqrt{V}} \sum_{r,k} \left[U_r(\vec{k}) c_{r,\vec{k}} + V_r(-\vec{k}) \tilde{c}_{r,-\vec{k}}^\dagger \right] e^{i\vec{k} \cdot \vec{x}}, \quad (5)$$

where U_r and V_r are given by

$$U_r(\vec{k}) = \begin{pmatrix} \cos \frac{\chi(\vec{k})}{2} \\ \vec{\sigma} \cdot \hat{k} \sin \frac{\chi(\vec{k})}{2} \end{pmatrix} u_r; \quad V_r(-\vec{k}) = \begin{pmatrix} -\vec{\sigma} \cdot \hat{k} \sin \frac{\chi(\vec{k})}{2} \\ \cos \frac{\chi(\vec{k})}{2} \end{pmatrix} v_r. \quad (6)$$

For free spinor fields, we have $\cos \chi(\vec{k}) = M/\epsilon(\vec{k})$, $\sin \chi(\vec{k}) = |\vec{k}|/\epsilon(\vec{k})$ with $\epsilon(\vec{k}) = \sqrt{\vec{k}^2 + M^2}$. However, we will deal with interacting fields so that we take the ansatz $\cos \chi(\vec{k}) = M^*/\epsilon^*(\vec{k})$, $\sin \chi(\vec{k}) = |\vec{k}^*|/\epsilon^*(\vec{k})$, with $\epsilon^*(\vec{k}) = \sqrt{\vec{k}^{*2} + M^{*2}(\vec{k})}$, where $\vec{k}^*$ and $M^*(\vec{k})$ are the effective momentum and effective mass respectively. The equal time anti-commutation conditions are $[c_{r,\vec{k}}, c_{s,\vec{k}'}^\dagger]_+ = \delta_{rs} \delta_{\vec{k},\vec{k}'} = [\tilde{c}_{r,\vec{k}}, \tilde{c}_{s,\vec{k}'}^\dagger]_+$. The vacuum $|0\rangle$ is defined through $c_{r,\vec{k}} |0\rangle = \tilde{c}_{r,\vec{k}}^\dagger |0\rangle = 0$; one-particle states are written $|\vec{k}, r\rangle = c_{r,\vec{k}}^\dagger |0\rangle$; two-particle and three-particle uncorrelated states are written, respectively as $|\vec{k}, r; \vec{k}', r'\rangle = c_{r,\vec{k}}^\dagger c_{r',\vec{k}'}^\dagger |0\rangle$, and $|\vec{k}, r; \vec{k}', r'; \vec{k}'', r''\rangle = c_{r,\vec{k}}^\dagger c_{r',\vec{k}'}^\dagger c_{r'',\vec{k}''}^\dagger |0\rangle$, and so on.

We now introduce the short range correlation through an unitary operator method. The correlated wave function is [16] $|\Psi\rangle = e^{iS} |\Phi\rangle$ where $|\Phi\rangle$ is a Slater determinant and S is, in general, a n -body Hermitian operator, splitting into a 2-body part, a 3-body part, etc.. Consider now the expectation value of H ,

$$E = \frac{\langle \Psi | H | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \frac{\langle \Phi | e^{-iS} H e^{iS} | \Phi \rangle}{\langle \Phi | \Phi \rangle}. \quad (7)$$

At this point we restrict ourselves to the two-body correlation diagrams shown in figure 1. Let us denote the two-body correlated wave function by

$$\overline{|\vec{k}, r; \vec{k}', r'\rangle} = e^{iS} |\vec{k}, r; \vec{k}', r'\rangle \approx f_{12} |\vec{k}, r; \vec{k}', r'\rangle \quad (8)$$

where f_{12} is the short range correlation factor, the so-called Jastrow factor [17]. For simplicity, we have written $f_{12} = f(\vec{r}_{12})$, $\vec{r}_{12} = \vec{r}_1 - \vec{r}_2$. We take the assumption that $f(r) = 1 - (\alpha + \beta r) e^{-\gamma r}$ where α , β and γ are parameters.

At this point some remarks on the choice of the Jastrow factor are appropriate. The important effect of the short range correlations is the replacement, in the expression for the ground-state energy, of the interaction matrix element $\langle \vec{k}, r; \vec{k}', r' | V_{12} | \vec{k}, r; \vec{k}', r' \rangle$ by $\overline{\langle \vec{k}, r; \vec{k}', r' | V_{12} + t_1 + t_2 | \vec{k}, r; \vec{k}', r' \rangle} - \langle \vec{k}, r; \vec{k}', r' | t_1 + t_2 | \vec{k}, r; \vec{k}', r' \rangle$, where t_i is the kinetic energy operator of particle i . As argued by Moszkowski [18] and Bethe [19], it is expected that the true ground-state wave function of the nucleus containing correlations coincide with the independent particle, or Hartree-Fock wave function, for interparticle distances $r \geq r_{heal}$, where $r_{heal} \approx 1$ fm is the so-called ‘‘healing distance’’. This behavior is a consequence of the constraints imposed by the Pauli Principle. Moreover, although in general the correlation factor $f(r)$ may depend on the isospin and spin quantum numbers of the two-body channel, we assume, for simplicity, that it is a plain, state independent Jastrow factor. A natural consequence of having the correlations introduced by an unitary operator is the occurrence of a normalization constraint on $f(r)$,

$$\int (f^2(r) - 1) d^3r = 0. \quad (9)$$

The correlated ground state energy becomes

$$\begin{aligned}
\mathcal{E} = & \frac{\nu}{\pi^2} \int_0^{k_F} k^2 dk [|k| \sin \chi(k) + M \cos \chi(k)] + \frac{\tilde{F}_\sigma(0)}{2} \rho_s^2 + \frac{\tilde{F}_\omega(0)}{2} \rho_B^2 \\
& + C_0 \rho_B \frac{\nu}{\pi^2} \int_0^{k_F} k^2 dk [|k| \sin \chi(k) + M \cos \chi(k)] \\
& - \frac{4}{(2\pi)^4} \int_0^{k_f} k^2 dk k'^2 dk' \left\{ [|k| \sin \chi(k) + 2 M \cos \chi(k)] I(k, k') + |k| \sin \chi(k') J(k, k') \right\} \\
& + \frac{1}{(2\pi)^4} \int_0^{k_f} k dk k' dk' \left[\sum_{i=\sigma, \omega} A_i(k, k') + \cos \chi(k) \cos \chi(k') \sum_{i=\sigma, \omega} B_i(k, k') + \sin \chi(k) \sin \chi(k') \sum_{i=\sigma, \omega} C_i(k, k') \right]
\end{aligned} \tag{10}$$

where, $C_0 = \int (f^2(r) - 1) d^3r$ so that, according to (9), $C_0 = 0$, and A_i, B_i, C_i, I and J are exchange integrals. In the above equation for the energy density, the first term results from the kinetic contribution, the second and third terms come respectively from the σ and ω direct contributions from the potential energy with correlations, the fourth and fifth ones from the direct and exchange correlation contribution from the kinetic energy, and the last one from the $\sigma + \omega$ exchange contributions from the potential energy with correlations. The angular integrals are given by $A_i(k, k') = B_i(k, k') = 2\pi g_i^2/4\pi \int_0^\pi d\cos\theta \tilde{F}_i(k, k', \cos\theta)$, $C_i(k, k') = 2\pi g_i^2/4\pi \int_0^\pi \cos\theta d\cos\theta \tilde{F}_i(k, k', \cos\theta)$, $I(k, k') = 2\pi \int_0^\pi d\cos\theta \tilde{C}_1(k, k', \cos\theta)$, and $J(k, k') = 2\pi \int_0^\pi \cos\theta d\cos\theta \tilde{C}_1(k, k', \cos\theta)$, where

$$\tilde{F}_i(\vec{k}, \vec{k}') = \int [f(r)V_\tau(r)f(r)] e^{i(\vec{k}-\vec{k}')\cdot\vec{r}} d\vec{r} \quad \text{and} \quad \tilde{C}_1(\vec{k}, \vec{k}') = \int (f^2(r) - 1) e^{i(\vec{k}-\vec{k}')\cdot\vec{r}} d\vec{r}. \tag{11}$$

More explicitly, the first terms of the above angle integrals read for $f(r) = 1$:

$$\begin{aligned}
A_\sigma &= g_\sigma^2 \theta_\sigma & B_\sigma &= g_\sigma^2 \theta_\sigma & C_\sigma &= -2g_\sigma^2 \phi_\sigma \\
A_\omega &= 2g_\omega^2 \theta_\omega & B_\omega &= -4g_\omega^2 \theta_\omega & C_\omega &= -4g_\omega^2 \phi_\omega \\
\theta_i(p, p') &= \log \left(\frac{(k+k')^2 + m_i^2}{(k-k')^2 + m_i^2} \right) & \phi_i(p, p') &= \frac{k^2 + k'^2 + m_i^2}{4kk'} \theta_i(p, p') - 1.
\end{aligned}$$

The baryon density and the scalar density are

$$\rho_B = \frac{2\nu}{(2\pi)^3} \int_0^{k_f} d\vec{k} = \frac{2\nu k_f^3}{6\pi^2}, \quad \rho_s = \frac{2\nu}{(2\pi)^3} \int_0^{k_f} \cos \chi(\vec{k}) d\vec{k}. \tag{12}$$

The parameters of the model have to be fixed. They are the couplings g_σ, g_ω , the meson masses, m_σ and m_ω and also three more parameters from the short range correlation function, α, β and γ . The couplings are chosen so as to satisfy the ground state properties of the nuclear matter and are given in table I. We choose $m_\sigma = 550$ MeV and take $m_\omega = 783$ MeV. The normalization condition (9) determines β . We fix α either by imposing the condition $f(0) = 0$, which appears to be a natural choice from our experience with the non-relativistic case, or by minimizing the energy. We choose γ so that the correct healing distance [18] is reproduced. If α is chosen to be 1, we assume a density independent parameter γ equal to 750 (HF+corr-IV). On the other hand, if we choose to determine α variationally, we assume that the parameter γ increases linearly with the Fermi momentum. The last choice is consistent with the idea that the healing distance decreases as k_F increases. Of course there are other possible choices for these parameters. The parameters we have used are tabulated in table I together with the compressibility K , the relative effective mass M^*/M , the kinetic energy $\mathcal{T}/\rho_B - M$, the direct and exchange parts of the potential energy ($\mathcal{V}_d/\rho_B$ and $\mathcal{V}_e/\rho_B$ respectively) with correlation and the correlation contribution to the kinetic energy $\mathcal{T}^C/\rho_B$, all computed at the saturation point. We next compute the range of the correlations, obviously related with the healing distance, also included in table I and defined as [19]:

$$R^2 = \frac{\int r^2 (f(r) - 1)^2 d^3r}{\int (f(r) - 1)^2 d^3r}. \tag{13}$$

In figure 2 we plot the binding energies as function of the density for the Hartree, HF, quark-meson-coupling model (QMC) [20], a non-linear Walecka model NL3 [21] and the 4 choices of the parameters in our calculation, as given in table I. From this figure one can see that in all cases that inclusion of correlations make the equation of state (EOS)

softer than Hartree or HF calculations. NL3 and QMC also provide softer EOS around nuclear matter saturation density but around two times saturation density, some of the EOS with correlations are softer than NL3. Correlations always tend to soften the EOS, except when $\alpha = 1$ is fixed. In this case, the EOS coincides with the curve obtained with the Hartree calculation.

In figure 3 we plot the individual contributions to the energy density in function of the density for the Hartree, HF and HF+correlations (set III). Notice that the correlation contribution is of the same order as the exchange contribution. Hence, it cannot be disregarded. In fact if we compare the coupling constants g_σ and g_ω obtained for the different calculations we conclude that the introduction of the correlations reduces the coupling constants. Correlation effects in the Hartree and HF calculations are taken into account by a correct choice of the coupling constants. However, the explicit introduction of correlations has other effects such as softening the EOS.

In conclusion, we have included two-body correlations in the $\sigma - \omega$ relativistic model. We have seen that they correspond to an important amount of the energy density and provide a softer EOS as compared with the HF approximation. Furthermore, the effect of the correlations is similar to the role played the non-linear terms in relativistic models or the 3-body forces in non-relativistic potential models. The inclusion of three-body correlations is currently under investigation.

This work was partially supported by CNPq (Brazil), CAPES (Brazil), GRICES (Portugal) under project 100/03, FEDER and FCT (Portugal) under the projects POCTI/FP/FNU/50326/2003 and POCTI/FIS/451/94.

-
- [1] M.H. MacGregor, A. R. Arndt, and R.M. Wright, Phys. Rev. **182**, 1714 (1969) and references therein.
 - [2] D.R. Entem and R. Machleidt, Phys. Lett B **524**, 93 (2001).
 - [3] R. Machleidt, Adv. Nucl. Phys. **19**, 189 (1989); R. Machleidt, Phys. Rev. **C 63**, 024001 (2001).
 - [4] R. Wiringa, R. Smith, and T. Ainsworth, Phys. Rev C **29**, 1207 (1984).
 - [5] F. Coester, S. Cohen, B.D. Day, and C.M. Vincent, Phys. Rev. **C 1**, 769 (1970); R. Brockmann and R. Machleidt, Phys. Rev. **C 42**, 1965 (1990).
 - [6] R.B. Wiringa, V. Fiks and A. Fabrocini, Phys. Rev. **C 38**, 1010 (1988); W. Zuo, A. Lejeune, U. Lombardo and J.-F. Mathiot, Nucl. Phys. **A 706**, 418 (2002).
 - [7] H.P. Dürr and E. Teller, Phys. Rev. **101**, 494 (1956); M. H. Johnson and E. Teller, Phys. Rev. **98**, 783 (1955); H.P. Dürr, Phys. Rev. **103**, 469 (1956).
 - [8] B.D. Serot and J.D. Walecka, Adv. Nucl. Phys. **16**, 1 (1986); J.D. Walecka, Ann. of Phys. **83**, 491 (1974); B.D. Serot, J.D. Walecka, Int. J. Mod. Phys. **E6**, 515 (1997).
 - [9] C.J. Horowitz and B.D. Serot, Nucl. Phys. **A399**, 529 (1983).
 - [10] A. Bouyssy, J.-F. Mathiot and N.V. Giai and S. Marcos, Phys. Rev. **C 36**, 380 (1987).
 - [11] F. Villars, "Proceedings of the International School of Physics, 'Enrico Fermi'-Course 23, (1961)." Academic Press, New York, 1963; J.S. Bell, " Lectures on the Many-Body Problem, First Bergen International School of Physics." Benjamin, New York, (1962); F. Coester and H. Kümmel, Nucl. Phys. **17** 477 (1960).
 - [12] J. da Providencia and C.M. Shakin, Ann. Phys.(NY) **30**, 95 (1964).
 - [13] K. Suzuki, R. Okamoto and H. Kumagai, Phys. Rep. **242** 181 (1994).
 - [14] H. Feldmeier, T. Neff, R. Roth, and J. Schnack, Nucl. Phys. **A 632**, 61 (1998); T. Neff and H. Feldmeier, Nucl. Phys. **713** 311 (2003).
 - [15] A. Mishra, P.K. Panda, S. Schramm, J. Reinhardt and W. Greiner, Phys. Rev. **C 56**, 1380 (1997).
 - [16] J. da Providencia and C. M. Shakin, Phys. Rev **C 4**, 1560 (1971); C. M. Shakin, Phys. Rev **C 4**, 684 (1971).
 - [17] R. Jastrow, Phys. Rev. **98**, 1479 (1955).
 - [18] S.A. Moszkowski and B.L. Scott, Ann. Phys. (N.Y.), **11**, 65 (1960).
 - [19] H. Bethe, Ann. Rev. Nucl. Sci. **21**, 93 (1971).
 - [20] P. A. M. Guichon, Phys. Lett. **B 200**, 235 (1988). K. Saito and A.W. Thomas, Phys. Lett. **B 327**, 9 (1994); P.K. Panda, A. Mishra, J.M. Eisenberg, W. Greiner, Phys. Rev. **C 56**, 3134 (1997).
 - [21] G. A. Lalazissis, J. König and P. Ring, Phys. Rev. **C 55**, 540 (1997).

TABLE I: Model parameters and ground state properties of nuclear matter at saturation density are given. These results were obtained with fixed: $M = 939$ MeV $m_\sigma = 550$ MeV, $m_\omega = 783$ MeV at $k_{F0} = 1.3 \text{ fm}^{-1}$ with binding energy $E_B = \varepsilon/\rho - M = -15.75$ MeV. We have used 4 choices of the parameters for the correlation. HF+corr-I corresponds to $600 + 200k_F/k_{F0}$, HF+corr-II to $700 + 200k_F/k_{F0}$, HF+corr-III to $600 + 300k_F/k_{F0}$ and HF+corr-IV corresponds to $\alpha = 1$ and γ fixed.

	g_σ	g_ω	α	β (MeV)	γ (MeV)	R fm	K (MeV)	M^*/M	$\mathcal{T}/\rho_B - M$ (MeV)	$\mathcal{V}_d/\rho_B$ (MeV)	$\mathcal{V}_e/\rho_B$ (MeV)	$\mathcal{T}^C/\rho_B$ (MeV)
Hartree	11.079	13.806					540	0.540	8.11	-23.86		
HF	10.432	12.223					585	0.515	5.87	-37.45	15.83	
HF+corr-I	4.326	2.431	13.474	-1867.295	800	0.338	523.	0.568	14.96	-96.81	26.77	39.33
HF+corr-II	4.359	2.4893	13.655	-2068.539	900	0.303	459.	0.608	15.8	-83.01	23.4	28.0
HF+corr-III	4.359	2.4893	13.655	-2068.539	900	0.303	448.	0.608	15.8	-83.01	23.4	28.0
HF+corr-IV	11.836	13.608	1.0	-244.678	750	0.318	573.	0.515	7.16	-45.17	17.94	4.31




FIG. 1: The kinetic and the potential contribution to the two-body cluster diagrams




FIG. 2: EoS for different parametrizations as defined in the table.




FIG. 3: Individual contributions for the energy density